


## 4.13

### Logarithmic Functions and Their Graphs

The inverse of an exponential function  $f(x) = b^x$  with base  $b$  is the logarithmic function with base  $b$ ,  $f^{-1}(x) = \log_b x$ .



The domain of the logarithm function is  $x > 0$ .

Domain of a logarithm function = range of the exponential function =  $(0, \infty)$  and the range of a logarithm function = domain of exponential function  $(-\infty, \infty)$ .

#### Changing Between Logarithmic and Exponential Form

If  $x > 0$  and  $0 < b \neq 1$ , then  $y = \log_b(x)$  (read as “ $y$  is the logarithm to the base  $b$  of  $x$ ”)

if and only if  $b^y = x$ .

#### Evaluating Logarithms

Examples:

b)  $\log_3 \sqrt{3} = \frac{1}{2}$ , because  $3^{\frac{1}{2}} = \sqrt{3}$

c)  $\log_5 \frac{1}{25} = -2$ , because  $5^{-2} = \frac{1}{5^2} = \frac{1}{25}$

d)  $\log_4 1 = 0$ , because  $4^0 = 1$

e)  $\log_7 7 = 1$ , because  $7^1 = 7$

### **Basic Properties of Logarithms**

For  $0 < b \neq 1$ ,  $x > 0$ , and any real number  $y$ ,

- $\log_b 1 = 0$  because  $b^0 = 1$ .
- $\log_b b = 1$  because  $b^1 = b$ .
- $\log_b b^y = y$  because  $b^y = b^y$ .
- $b^{\log_b x} = x$  because  $\log_b x = \log_b x$ .

### **Evaluating Logarithmic and Exponential Expressions**

**Examples:**

a)  $\log_2 8 = \log_2 2^3 = 3$

b)  $\log_3 \sqrt{3} = \log_3 3^{\frac{1}{2}} = \frac{1}{2}$

c)  $6^{\log_6 11} = 11$

### **Basic Properties of Common Logarithms**

**Let  $x$  and  $y$  be real numbers with  $x > 0$ .**

$\log 1 = 0$  because  $10^0 = 1$

**$\log 10 = 1$**  because  $10^1 = 10$

**$10^{\log x} = x$**  because  $\log x = \log x$

**$\log 10^y = y$**  because  $10^y = 10^y$

## Evaluating Logarithmic & Exponential Expressions—Base 10

### Examples:

a)  $\log 100 = \log_{10} 100 = 2$ , *because*  $10^2 = 100$

b)  $\log \sqrt[5]{10} = \log 10^{\frac{1}{5}} = \frac{1}{5}$

c)  $\log \frac{1}{1000} = \log \frac{1}{10^3} = \log 10^{-3} = -3$

d)  $10^{\log 6} = 6$

## Evaluating Common Logarithms with a Calculator

### Examples:

a)  $\log 34.5 = 1.537\dots$ , *because*  $10^{1.537\dots} = 34.5$

b)  $\log 0.43 = -0.366\dots$ , *because*  $10^{-0.366\dots} = 0.43$

c)  $\log(-3)$  is undefined because there is no real number  $y$  such that  $10^y = -3$ .

## Solving Simple Logarithmic Equations

**Examples:** Solve each equation by changing it to exponential form.

a)  $\log x = 3$

Change to exponential form,  $x = 10^3 = 1000$

b)  $\log_2 x = 5$

Change to exponential form,  $x = 2^5 = 32$

## Basic Properties of Natural Logarithms

Let  $x$  and  $y$  be real numbers with  $x > 0$

$\ln 1 = 0$  because  $e^0 = 1$ .

$\ln e = 1$  because  $e^1 = e$ .

$e^{\ln x} = x$  because  $\ln x = \ln x$ .

$\ln e^y = y$  because  $e^y = e^y$ .

## Evaluating Logarithmic & Exponential Expressions Base – e

Examples:

a)  $\ln \sqrt{e} = \log_e \sqrt{e} = \frac{1}{2}$ , because  $e^{\frac{1}{2}} = \sqrt{e}$

b)  $\ln e^5 = \log_e e^5 = 5$

c)  $e^{\ln 4} = 4$

## Evaluating Natural Logarithms with a Calculator

Examples:

a)  $\ln 23.5 = 3.157 \dots$ , because  $e^{3.157 \dots} = 23.5$

b)  $\ln 0.48 = -0.733 \dots$ , because  $e^{-0.733 \dots} = 0.48$

c)  $\ln (-5)$  is undefined because there is no real number  $y$  such that  $e^y = -5$ .

## The Natural Logarithmic Function

$$f(x) = \ln x$$

Domain:  $(0, \infty)$

Range: All reals

Continuous on  $(0, \infty)$

Increasing on  $(0, \infty)$

No symmetry

Not bounded above or below

No local extrema

No horizontal asymptotes

Vertical asymptote:  $x = 0$

End Behavior:  $\lim_{x \rightarrow \infty} \ln x = \infty$

(Do examples of graphing.)

